



## GS-441524 INFORMATION SHEET

**GS-441524 is an antiviral drug which has been shown to be highly effective in treating Feline Infectious Peritonitis (FIP). Without treatment, FIP is 100% fatal.**

### ADMINISTERING THIS MEDICATION

This liquid oral suspension can be given directly into your cat's mouth via the syringe provided, or mixed into a small amount of food or a lickable treat. It is recommended to give this medication on an empty stomach, and to not feed a large meal for about an hour afterwards.

Your veterinarian may direct you to give your cat this medication either once or twice daily. It is very important to make sure that your cat receives their prescribed dose(s) every day. If you miss a dose, give it as soon as you remember, and then continue with the regular dosing schedule. However, if it is almost time for the next dose, skip the missed dose and resume the regular dosing schedule. Failure to give your cat their medication every day as directed may lead to treatment failure.

12 weeks of antiviral treatment (such as with GS- 441524) is the current recommendation for FIP treatment. Do not stop treatment without consulting with your veterinarian. Stopping treatment prematurely can cause your cat to relapse and become sick again.

If you are having difficulty administering this medication, consult your veterinarian or pharmacist for advice on administration techniques. For online tips on administering oral liquid suspensions to your cat, please check out the International Cat Care group's tips on medicating a cat: <https://icatcare.org/articles/how-to-give-a-cat-medication>

### FORMULATION

**Dosage Form:** Oral Suspension

**Available Volumes:** 30 mL and above qualifies for free shipping, but you may order any volume you prefer

**Strength:** 50mg/mL

**Available Flavours:** Fish or Chicken





## STORAGE AND HANDLING

GS-441524 oral suspension should be stored at room temperature.

There are no specific precautions for handling this medication unless you are allergic. Always wash your hands after handling or administering any medication.

## MONITORING

If your cat experiences new or worsening symptoms, consult your veterinarian, as a dosage adjustment or additional supportive care may be necessary.

It is very important to monitor your cat's weight (weekly) during treatment so that the dose can be adjusted for weight gain. Failure to adjust the dose for weight gain is a leading cause of treatment failure.

